

## Exploring Hilbert space: Accurate characterization of quantum information

A. G. White,<sup>1,\*</sup> D. F. V. James,<sup>2</sup> W. J. Munro,<sup>1,3</sup> and P. G. Kwiat<sup>2,4</sup>

<sup>1</sup>Department of Physics, University of Queensland, Brisbane, Queensland 4072, Australia

<sup>2</sup>Theory Division, T-4 and Physics Division, P-23, Los Alamos National Laboratory, Los Alamos, New Mexico 87545

<sup>3</sup>Maths, Cryptography and Security Group, Hewlett-Packard Laboratories, Bristol BS34 8QZ, United Kingdom

<sup>4</sup>Department of Physics, University of Illinois, Urbana-Champaign, Illinois 61801

(Received 2 July 2001; published 4 December 2001)

We report the creation of a wide range of quantum states with controllable degrees of entanglement and entropy using an optical two-qubit source based on spontaneous parametric down-conversion. The states are characterized using measures of entanglement and entropy determined from tomographically determined density matrices. The tangle-entropy plane is introduced as a graphical representation of these states, and the theoretic upper bound for the maximum amount of entanglement possible for a given entropy is presented. Such a combination of general quantum state creation and accurate characterization is an essential prerequisite for quantum device development.

DOI: 10.1103/PhysRevA.65.012301

PACS number(s): 03.67.-a, 03.65.Ud, 03.65.Wj, 03.65.Yz

Quantum information (QI) — the application of quantum mechanics to problems in information science such as computation and communication — has led to a renewed interest in fundamental aspects of quantum mechanics. In particular much attention has been focused on the role of *entanglement*, the nonclassical correlation between separate quantum systems, particularly, between two-level systems or *qubits* (quantum bits). Entanglement, along with the degree of order, or *purity*, determines the utility of a given system for realizing various QI protocols. A key goal of QI is the experimental realization of complex quantum algorithms, e.g., Shor’s algorithm, which allows efficient factoring of composite integers [1]: recent research indicates that while entanglement is necessary to execute Shor’s algorithm, pure states are not [2].

There is currently a global effort to manufacture two-qubit gates, since any quantum algorithm can be implemented by a combination of single-qubit rotations and such gates (which produce the necessary entanglement) [3]. These are fully characterized only when both the gate states and its dynamics have been accurately measured, which requires a tunable source of two-qubit quantum states and a method of completely measuring the output states [4]. No system to date has fulfilled these criteria. Here we report an optical two-qubit source that produces a wide range of quantum states with controllable degrees of purity and entanglement, and fully characterize these states by quantum tomography. This source is also suitable for exploring alternative paradigms: (1) where quantum algorithms are implemented via single-qubit rotations, Bell-state measurements, and a predetermined set of entangled states (that may or may not need to be pure) [5]; (2) scalable linear-optics quantum computation, where entanglement occurs as a result of measurement [6].

Quantum states of  $N$  qubits can be represented by a vector existing in a  $2^N$ -dimensional Hilbert space. This is the

“space of possibilities,” and represents all possible physical combinations of qubits for a system. To date the states generated in QI experiments have clustered around two distinct limits in Hilbert space: (1) highly entangled systems with high order [7–11]; (2) completely unentangled systems with very little order [12]. The lack of entanglement in the latter case [13] has raised the question of what properties are actually required for quantum information protocols, and highlights that to date, the “domain of mixed states between these two extremes [pure vs completely mixed] is incredibly big and largely unexplored” [14]. We experimentally explore this unmapped region, and introduce a theoretical upper bound for the maximum amount of entanglement possible for a given purity. A variety of measures exist for quantifying the degrees of disorder and entanglement, all of which are functions of the system density matrix. For our experimental system, the density matrix can now be obtained via quantum tomography [15], allowing these measures to be applied. In this paper we will use the *tangle*,  $T$ , to quantify the degree of entanglement, and the *linear entropy*,  $S_L$ , to quantify the degree of disorder [16–19].

We obtain our quantum states via spontaneous down-conversion, where a pump photon passed through a nonlinear crystal is converted into a pair of lower-energy photons. We use the polarization state of the single photons as our qubits, and measure in coincidence, thus obtaining the reduced density matrix (it only describes the polarization component of the state) of the two-photon contribution [20,21]. Figure 1 is a schematic of the experimental system, a detailed descrip-




FIG. 1. Experimental setup for quantum-state synthesis.

\*Author to whom correspondence and requests for materials should be addressed. Email address: andrew@physics.uq.edu.au




FIG. 2. States obtained by spatially based polarization decoherence. (a) (i) Input state  $(|HH\rangle + |VV\rangle)/\sqrt{2}$ ; (ii),(iii) states after passing through BBO decoherers in one and both arms, respectively. States obtained by temporally based polarization decoherence. (b) The state,  $(|HH\rangle + |VV\rangle)/\sqrt{2}$ , after passing through the state selector ( $\theta_1=0$ ,  $\theta_2$  set as shown) and the quartz decoherers set as described in the text. (c) As for (b), but with the nonmaximally entangled initial state  $0.96|HH\rangle + 0.29|VV\rangle$ . Only the real components of the density matrices are shown, the imaginary components being at the few percent level or less.

tion is given elsewhere [22]. Briefly, the beta-barium-borate (BBO) crystals produce pure-state pairs of photons that can be tuned between the separable and maximally entangled limits by adjusting the pump polarization [15]. The parity and phase of the entangled states are selected via the “state selector” half-wave plates, and the photons are analyzed using adjustable quarter- and half-wave plates (HWP) and polarizing beamsplitters, which enable polarization analysis in any basis. (For tomography, 16 different coincidence bases are required [15,23]). The photons are passed via suitable optics to single-photon counters, whose outputs are recorded in coincidence.

To change the entropy it is necessary to introduce decoherence into the polarization degree of freedom, which can be done either spatially or temporally. Decoherence can occur when the phase,  $\phi$ , of the entangled state (e.g.,  $|HH\rangle + e^{i\phi}|VV\rangle$ ) varies rapidly over a small spatial extent, i.e., smaller than the collection apertures. We achieved this by the introduction of BBO crystals (3 mm thick) into the down-conversion beams, cut so that their optic axes were at an

angle of  $49^\circ$  to the beam. These introduce a highly direction-dependent phase shift in the down-converted photons: due to the intrinsic spread of photon momentum in down-conversion, and the high birefringence of BBO, after the decoherer the phase of the entanglement is very finely fringed compared to the collection aperture. Figure 2(a) shows that with BBO in only one arm (optic axis at  $45^\circ$ ), the resultant state is partially mixed and completely unentangled,  $(S_L, T) = (0.66 \pm 0.03, 0.00 \pm 0.00)$ ; adding a BBO crystal to the remaining arm, (optic axis at  $0^\circ$ ) generates a fully mixed state,  $(S_L, T) = (1.00 \pm 0.01, 0.00 \pm 0.00)$ .

As spatially based decoherence completely destroys entanglement, it is not a suitable technique for exploring Hilbert space. In contrast, temporally based polarization decoherence allows entanglement to survive. It is achieved by imposing a large relative phase delay, longer than the coherence length of the light, between two orthogonal polarizations. In practice this is realized by introducing a 10-mm-thick quartz crystal in each arm, with optic axes vertical and perpendicular to the beam. The detected photons have a co-




FIG. 3. Tangle vs linear entropy for two qubits. Black curve: Werner states. Data points are calculated tangle and linear entropy from a range of measured density matrices. The gray region indicates physically impossible combinations of  $T$  and  $S_L$ ; maximal states [Eq. (1)] lie at the boundary of this region.

herence length of 140 wavelengths (set by the 5-nm interference filters), 10 mm of quartz delays the phase velocity of the horizontally polarized light by this amount (relative to the vertical). Viewed differently, the quartz entangles the phase to the photon frequency, which is then traced over [24,25]. Thus, a single photon linearly polarized at  $45^\circ$  would exit the quartz crystal strongly depolarized, i.e., in a mixed state; similarly, a pair of photons, each at  $45^\circ$ , would exit two such crystals in a mixed state. With entangled states, however, the situation is more subtle. Certain kinds of entangled states are immune to collective decoherence, whilst others exhibit strong decoherence—the former comprise *decoherence-free subspaces* [26]. Due to the energy entanglement of the photons, and the alignment of our decoherers, in our system the state  $(|HH\rangle + |VV\rangle)/\sqrt{2}$  is decoherence-free [24,27]. The function of the state selector is to continuously tune this state towards another maximally entangled state, one that is not decoherence-free [e.g.,  $(|HV\rangle + |VH\rangle)/\sqrt{2}$ ].

Figures 2(b) and 2(c) show a range of density matrices generated via this method. In Fig. 2(b), selector wave plate HWP1 was fixed at  $0^\circ$  and HWP2 varied by the angle indicated, Fig. 2(c) shows a similar series, except this time starting with a nonmaximally entangled state. There are 16 parameters in the density matrix, too many for easy assimilation. To see how much of Hilbert space we are accessing with these states, we use the tangle and linear entropy measures to construct a characteristic plane as a succinct, compact way of representing the salient features of a quantum state. In this plane (Fig. 3), a pure, unentangled state lies at the origin  $(S_L, T) = (0, 0)$ ; a pure, maximally entangled state in one corner  $(S_L, T) = (0, 1)$ ; and a maximally mixed, unentangled state in the corner diagonally opposite  $(S_L, T) = (1, 0)$ . A maximally entangled, maximally mixed state  $(S_L, T) = (1, 1)$  is obviously impossible. As indicated above, previous QI experiments have generated states either near the tangle axis ( $S_L \sim 0, 0 \leq T \leq 1$ ) (cavity QED, ion and photon experiments) or at the maximally mixed point ( $S_L \sim 1 - 10^{-4}$  to  $1 - 10^{-6}$ ,  $T = 0$ ) (high-temperature nuclear

magnetic-resonance experiments). In Fig. 3, we plot the linear entropy and tangle values determined from a range of our measured density matrices, including those shown in Fig. 2 and from [28]. Two sources of experimental uncertainty were considered, statistical uncertainties ( $\sqrt{N}$ , where  $N$  is the count), which range from 1–4% for our count rates, and settings uncertainties, due to the fact that the analyzers can only be set with an accuracy of  $\pm 0.25^\circ$ . The combination of these effects led to the uncertainties as shown, a full derivation of their calculation is lengthy and given elsewhere [23]. The heavy black line in Fig. 3 represents the  $(S, T)$  values of Werner states  $\hat{\rho}_W = \lambda \hat{\rho}_{mix} + (1 - \lambda) \hat{\rho}_{ent}$ , where  $\hat{\rho}_{mix}$  is a maximally mixed state,  $\hat{\rho}_{ent}$  is a pure maximally entangled state, and  $0 < \lambda < 1$  [29]. A wide array of states were created, up to and lying on the Werner border. Interestingly, for linear entropies less than  $8/9$ , there exist states with greater tangle than Werner states, the largest of which, the *maximal* states, lie at the boundary of the gray region in Fig. 3. The density matrix for these states has the form [30]

$$\hat{\rho} = \begin{pmatrix} D & 0 & 0 & \sqrt{T}/2 \\ 0 & 1 - 2D & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \sqrt{T}/2 & 0 & 0 & D \end{pmatrix}, \quad (1)$$

where  $D = \sqrt{T}/2$  when  $\sqrt{T} \geq 2/3$ , and  $D = 1/3$  when  $\sqrt{T} < 2/3$ . Only states with tangle and linear entropy that fall on or under this boundary are physically realizable. Using the current scheme of two decohering crystals, it is not possible to create states that lie between the Werner and maximal boundaries. We are currently investigating a method to realize generalized arbitrary quantum-state synthesis, which will allow generation of states with any allowed combination of entropy and tangle.

In any experimental system mixture is inevitable—our source enables experimental investigation of decoherence-induced effects and issues including entanglement purification [31], distillation [28,32], concentration [33], decoherence-free subspaces [26,27], and protocols that *require* decoherence [34]. Since decoherence is controllable in our system, it can be used as a testbed for controlled exploration of the effect of intrinsic, uncontrollable, decoherence in other architectures (e.g., decoherence in a solid-state two-qubit gate [35]).

Mixed entangled states also have fundamental ramifications. Entanglement, as defined by Schrödinger, is essentially a pure-state concept (resting as it does on the issue of separability) and is “. . . the quintessential feature of quantum mechanics, the one that enforces its entire departure from classical lines of thought” [36]. Is this indeed the case for mixed entangled states, or is there some other, perhaps operational, characteristic that would better define the boundary between quantum and classical mechanics? For example, we can make states that are mixed and nonseparable and yet do not violate a Bell’s inequality—are these “truly” entangled? Distilling these states makes states that are more mixed and more entangled and so that they now violate a Bell’s inequal-

ity [28]—was this entanglement really “hidden”? Using the tangle criteria, the answer is straightforward, the states were always entangled and the distillation simply moved their position on the tangle-entropy plane across the Bell boundary [37]. Yet, states that do not violate Bell’s inequality can be described by a hidden local-variable model, suggesting that some entangled states are classical, in violation of Schrödinger’s precept. The question of what significant physical

differences, if any, exist between these various mixed entangled states remains open.

D.F.V.J. wishes to thank the University of Queensland for their hospitality. This work was supported in part by the Australian Research Council (ARC), the U.S. National Security Agency (NSA), and the U.S. Advanced Research and Development Activity (ARDA).

- 
- [1] P. W. Shor, in *Proceedings of 35th Annual Symposium on Found. Computer Science*, edited by S. Goldwasser (IEEE, Los Alamitos, CA, 1994), p. 116.
- [2] S. Parker and M. B. Plenio, e-print quant-ph/0102136.
- [3] D. P. DiVincenzo, *Phys. Rev. A* **51**, 1015 (1995).
- [4] I. L. Chuang and M. A. Nielsen, *J. Mod. Opt.* **44**, 2455 (1997).
- [5] D. Gottesman and I. L. Chuang, *Nature (London)* **402**, 390 (1999).
- [6] E. Knill, R. Laflamme, and G. J. Milburn, *Nature (London)* **409**, 46 (2001).
- [7] Q. A. Turchette, C. J. Hood, W. Lange, H. Mabuchi, and H. J. Kimble, *Phys. Rev. Lett.* **75**, 4710 (1995).
- [8] K. Mattle, H. Weinfurter, P. G. Kwiat, and A. Zeilinger, *Phys. Rev. Lett.* **76**, 4656 (1996).
- [9] D. Bouwmeester *et al.*, *Nature (London)* **390**, 575 (1997).
- [10] Q. A. Turchette *et al.*, *Phys. Rev. Lett.* **81**, 3631 (1998).
- [11] D. S. Naik, C. G. Peterson, A. G. White, A. J. Berglund, and P. G. Kwiat, *Phys. Rev. Lett.* **84**, 4733 (2000).
- [12] I. L. Chuang, N. Gershenfeld, and M. Kubinec, *Phys. Rev. Lett.* **80**, 3408 (1998).
- [13] S. L. Braunstein *et al.*, *Phys. Rev. Lett.* **83**, 1054 (1999).
- [14] H. J. Kimble, *Phys. Today* **53** (1), 22 (2001).
- [15] A. G. White, D. F. V. James, P. H. Eberhard, and P. G. Kwiat, *Phys. Rev. Lett.* **83**, 3103 (1999).
- [16] The tangle,  $T$ , is directly related to the entanglement of formation,  $E$ , which is regarded as the canonical measure of entanglement [17,18]. (Specifically,  $E = h[(1 + \sqrt{1-T})/2]$ , where  $h(x) = -x \log_2 x - (1-x) \log_2 (1-x)$ .) For our two-qubit system we use a normalized form of the linear entropy,  $S_L = 4(1-P)/3$ , where the state purity is  $P = \text{Tr}(\rho^2)$  [19]. Note that  $0 \leq S_L, T \leq 1$ .
- [17] W. K. Wootters, *Phys. Rev. Lett.* **80**, 2245 (1998).
- [18] V. Coffman, J. Kundu, and W. K. Wootters, *Phys. Rev. A* **61**, 052306 (2000).
- [19] S. Bose and V. Vedral, *Phys. Rev. A* **61**, 040101(R) (2000).
- [20] By looking in coincidence, we restrict ourselves to the two- (and more)-photon section of the Hilbert space, i.e., we can disregard the vacuum component. Two-photon states measured in this manner have been shown to near-maximally violate Bell’s inequality [21], indicating that states are near-pure and near-maximally entangled, i.e., strongly nonclassical.
- [21] G. Weihs, T. Jennewein, C. Simon, H. Weinfurter, and A. Zeilinger, *Phys. Rev. Lett.* **81**, 5039 (1998).
- [22] P. G. Kwiat, E. Waks, A. G. White, I. Appelbaum, and P. H. Eberhard, *Phys. Rev. A* **60**, R773 (1999).
- [23] D. F. V. James, P. G. Kwiat, W. J. Munro, and A. G. White, *Phys. Rev. A* **64**, 052312 (2001).
- [24] A. J. Berglund, e-print quant-ph/0010001.
- [25] We stress that this principle—tracing over another degree of freedom in a tightly controlled way to induce decoherence—is applicable well beyond our specific experimental system.
- [26] D. A. Lidar, I. L. Chuang, and K. B. Whaley, *Phys. Rev. Lett.* **81**, 2594 (1998).
- [27] P. G. Kwiat, A. J. Berglund, J. Altepeter, and A. G. White, *Science* **290**, 498 (2000).
- [28] P. G. Kwiat, S. Barraza-Lopez, A. Stefanov, and N. Gisin, *Nature (London)* **409**, 1014 (2001).
- [29] R. F. Werner, *Phys. Rev. A* **40**, 4277 (1989).
- [30] W. J. Munro, D. F. V. James, A. G. White, and P. G. Kwiat, *Phys. Rev. A* **64**, 030302(R) (2001).
- [31] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, *Phys. Rev. A* **54**, 3824 (1996).
- [32] V. Vedral, M. B. Plenio, M. A. Rippin, and P. L. Knight, *Phys. Rev. Lett.* **78**, 2275 (1997).
- [33] R. T. Thew and W. J. Munro, *Phys. Rev. A* **63**, 030302(R) (2001).
- [34] R. Cleve, D. Gottesman, and H.-K. Lo, *Phys. Rev. Lett.* **82**, 648 (1999).
- [35] B. Kane, *Nature (London)* **393**, 133 (1998).
- [36] E. Schrödinger, *Proc. Cambridge Philos. Soc.* **31**, 555 (1935).
- [37] W. J. Munro, K. Nemoto, and A. G. White, *J. Mod. Opt.* **48**, 1239 (2001).